

### Cattle-Lac Liquid Feed 35

Crude Protein, not less than (This includes not more than 25% equivalent protein from non-protein nitrogen)	35.00%	Zinc (Zn), not less than	180.00ppm
Crude Fat, not less than	0.75%	Vitamin A, not less than	25,000 I.U./lb
Moisture, not more than	40.00%	Vitamin D3, not less than	6,500 I.U./lb
Phosphorus (P), not less than	0.50%	Vitamin E, not less than	20.00 I.U./lb
Magnesium (Mg), not less than	0.20%	Thiamine (B1), not less than	0.30 mg/lb
Potassium (K), not less than	1.00%	Riboflavin (B2), not less than	0.67 mg/lb
Cobalt (Co), not less than	0.56ppm	Niacin, not less than	9.70 mg/lb
Copper (Cu), not less than	30.00ppm	Vitamin B6 (Pyridoxine), not less than	0.80 mg/lb
Iodine (I), not less than	2.50ppm	d - Pantothenic Acid, not less than	2.75 mg/lb
Iron (Fe), not less than	140.00ppm	Biotin, not less than	0.06 mg/lb
Manganese (Mn), not less than	42.00ppm	Folic Acid, not less than	0.034 mg/lb
Selenium (Se), not less than	0.40ppm	Choline, not less than	290.00 mg/lb

#### Ingredients:

Condensed Corn Distillers Solubles, Cane Molasses ("Black Strap"), Fish Solubles, Hemicellulose Extract, Urea, Concentrated Separated By-products, Phosphoric Acid, Sulfuric Acid, Vitamin A Supplement, D-Activated Plant Sterol (source of Vitamin D3), D1-Alpha-Tocopheryl Acetate (source of Vitamin E), Copper Sulfate, Cobalt Sulfate, Manganese Sulfate, Zinc Sulfate, Ethylenediamine Dihydrochloride (source of Iodine).

#### Feeding Directions:

Cattle-Lac Liquid 35 should be fed free choice in lick wheel feeders to RUMINANTS ONLY.

Place feeder away from where animals congregate (loafing areas).

One wheel should be available for each 25 head.

Cattle should consume from 1-3 pounds daily depending on supplementary feed, and quality and quantity of roughage.

Always provide adequate quantities of roughage, water and minerals at all times.