

Pasture Booster 32

Guaranteed Analysis

Crude Protein, not less than (This includes not more than 23.50% equivalent protein from non-protein nitrogen)	32.00%	Iodine (I), not less than	2.50ppm
Crude Fat, not less than	0.12%	Iron (Fe), not less than	120.00ppm
Moisture, not more than	40.00%	Manganese (Mn), not less than	50.00ppm
Calcium (Ca), not less than	0.20%	Selenium (Se), not less than	0.40ppm
Calcium (Ca), not more than	0.60%	Zinc (Zn), not less than	170.00ppm
Phosphorus (P), not less than	0.15%	Vitamin A, not less than	15,400 I.U./lb
Sulfur (S), not less than	0.60%	Vitamin D3, not less than	3,855 I.U./lb
Cobalt (Co), not less than	0.50ppm	Vitamin E, not less than	3.80 I.U./lb
Copper (Cu), not less than	20.00ppm		

Ingredients:

Condensed Corn Distillers Solubles, Cane Molasses ("Black Strap"), Fish Solubles, Hemicellulose Extract, Urea, Concentrated Separated By-products, Phosphoric Acid, Sulfuric Acid, Vitamin A Supplement, D-Activated Plant Sterol (source of Vitamin D3), D1-Alpha-Tocopheryl Acetate (source of Vitamin E), Copper Sulfate, Cobalt Sulfate, Manganese Sulfate, Zinc Sulfate, Ethylenediamine Dihydrochloride (source of Iodine).

Feeding Directions:

Pasture Booster 32 should be fed free choice in lick wheel feeders to RUMINANTS ONLY.

Place feeder away from where animals congregate (loafing areas).

One wheel should be available for each 25 head.

Cattle should consume from 1-3 pounds daily depending on supplementary feed, and quality and quantity of roughage.

Always provide adequate quantities of roughage, water and minerals at all times.